

**Table E-11. Age, BMI, and smoking status for trials comparing antidepressant with placebo**

<b>Study</b>	<b>Age, Mean (SD)</b>	<b>Range</b>	<b>BMI, Mean (SD)</b>	<b>Current</b>	<b>Former</b>	<b>Never</b>
Suvanto-Luukkonen 2005	54.0	45 to 66		24.8		
Evans 2005	52.2 (5.5)			30.2		
Kerwin 2007	52.5 (5.1)			14.9		
Kalay 2007	52.6 (4.9)		28.0 (4.2)	22		
Speroff 2008	53.5 (4.8)	37 to 78	27.0 (4.6)			
Soares 2008	56.3 (2.7)					
Archer 2009	53.7 (5.0)		27.1 (4.6)			
Archer 2009	53.4 (4.8)	29 to 71	27.9 (5.0)			
Kornstein 2010	52.3 (6.3)	40 to 70				
Soares 2010	56.0 (6.0)					
Freeman 2011	53.9 (4.1)		29.1 (6.5)	22.9	28.8	48.3
Bouchard 2012	54.0 (4.5)	40 to 66	26.0 (4.0)			
Pinkerton 2012	54.0 (5.0)	45 to 71	26.5 (4.0)			
Simon (Study 2) 2013		40 to 74	27.5			
Pinkerton 2013	54.0 (5.0)	43 to 77	26.7 (4.0)			
Simon (Study 1) 2013		40 to 79	28.7			

BMI: body mass index; (c): data came from posted results on the clinical trial registry; SD: standard deviation