

Appendix Table F101. Clinical outcomes after magnetic stimulation compared to no active treatment (results from RCTs pooled with random effects models)

Outcome	Reference	Active events/ randomized	Control events/ randomized	Relative risk (95% CII)	Weight, %	Absolute risk difference (95% CI)	Weight %	Inclusion of mixed UI
Continence	Fujishiro, 2000 ⁵²⁹	4/31	1/31	4.00 (0.47; 33.79)	4	0.10 (- 0.04; 0.23)	51	No
Continence	Gilling, 2009 ⁵³⁴	6/35	3/35	2.00 (0.54; 7.37)	11	0.09 (- 0.07; 0.24)	38	No
Continence	But, 2005 ⁵⁰⁹	17/23	11/16	1.08 (0.71; 1.62)	85	0.05 (- 0.24; 0.34)	11	Yes
Pooled		27/89	15/82	1.22 (0.78; 1.88)	100	0.09 (- 0.01; 0.18)	100	
Heterogeneity p value, I squared				0.35	4.00	0.96	0.00	
Improved UI	But, 2003 ⁵⁰⁸	7/30	1/22	5.13 (0.68; 38.77)	6	0.19 (0.01; 0.36)	46	Not reported
Improved UI	But, 2005 ⁵⁰⁹	11/26	3/13	1.83 (0.62; 5.45)	19	0.19 (- 0.11; 0.49)	21	Yes
Improved UI	Fujishiro, 2000 ⁵²⁹	23/31	10/31	2.30 (1.33; 3.99)	75	0.42 (0.19; 0.64)	33	No
Pooled		41/87	14/66	2.30 (1.43; 3.71)	100	0.27 (0.11; 0.42)	100	
Heterogeneity p value, I squared				0.68	0.00	0.25	27.90	