

Appendix Table F103. Scoring of quality of life after magnetic stimulation compared to no active treatment (results from RCTs)

Reference	Active	Definition of quality of life	Randomized active/control	Active mean/standard deviation	Control mean/standard deviation	Mean difference (95% CI)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	General health perception, 1 week (T2)	10/10	37.50/13.10	42.50/16.80	-5.00 (-18.20; 8.20)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Incontinence impact, 1 week (T2)	10/10	39.90/26.20	56.60/22.40	-16.70 (-38.06; 4.66)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Role limitation, 1 week (T2)	10/10	33.30/30.40	33.30/22.20	0.00 (-23.33; 23.33)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Physical limitation, 1 week (T2)	10/10	43.20/27.40	46.60/24.60	-3.40 (-26.22; 19.42)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Social limitation, 1 week (T2)	10/10	14.90/19.50	32.10/21.20	-17.20 (-35.05; 0.65)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Personal relationships, 1 week (T2)	10/10	6.60/11.60	31.60/39.60	-25.00 (-50.58; 0.58)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Emotions, 1 week (T2)	10/10	41.00/29.10	42.10/29.00	-1.10 (-26.56; 24.36)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Sleep/energy, 1 week (T2)	10/10	29.90/20.40	19.90/13.10	10.00 (-5.03; 25.03)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	SEAPI-QMM, 1 week (T2)	10/10	1.70/0.80	1.80/0.60	-0.10 (-0.72; 0.52)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	General health perception, 1 month (T3)	10/10	52.30/25.90	57.50/28.90	-5.20 (-29.25; 18.85)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Incontinence impact, 1 month (T3)	10/10	49.60/22.20	64.90/16.50	-15.30 (-32.44; 1.84)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Role limitation, 1 month (T3)	10/10	39.90/29.60	53.30/23.30	-13.40 (-36.75; 9.95)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Physical limitation, 1 month (T3)	10/10	47.90/28.80	58.20/26.30	-10.30 (-34.47; 13.87)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Social limitation, 1 month (T3)	10/10	27.20/33.00	44.40/28.60	-17.20 (-44.27; 9.87)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Personal relationships, 1 month (T3)	10/10	13.80/14.60	34.90/34.60	-21.10 (-44.38; 2.18)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Emotions, 1 month (T3)	10/10	46.30/30.90	48.80/35.10	-2.50 (-31.48; 26.48)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	Sleep/energy, 1 month (T3)	10/10	29.90/17.20	33.30/13.60	-3.40 (-16.99; 10.19)
Manganotti, 2007 ⁵⁶⁵	Active magnetic stimulation	SEAPI-QMM, 1 month (T3)	10/10	2.30/0.80	2.10/0.30	0.20 (-0.33; 0.73)

**Appendix Table F103. Scoring of quality of life after magnetic stimulation compared to no active treatment (results from RCTs)
(continued)**

Reference	Active	Definition of quality of life	Randomized active/control	Active mean/standard deviation	Control mean/standard deviation	Mean difference (95% CI)
Gilling, 2009 ^{5,34}	Electromagnetic stimulation	I-QOL score	35/35	71.20/3.30	67.30/4.40	3.90 (2.08; 5.72)
Gilling, 2009 ^{5,34}	Electromagnetic stimulation	KHQ score	35/35	6.90/0.70	8.60/1.00	-1.70 (-2.10; -1.30)
Gilling, 2009 ^{5,34}	Electromagnetic stimulation	I-QOL score at 6 months of followup	35/35	73.60/3.00	68.90/4.50	4.70 (2.91; 6.49)
Gilling, 2009 ^{5,34}	Electromagnetic stimulation	KHQ score at 6 months of followup	35/35	7.70/0.70	8.50/1.00	-0.80 (-1.20; -0.40)