

Appendix Table F155. Comparative effectiveness of intravaginal electrical stimulation and trospium hydrochloride in women with overactive bladder syndrome (results from individual randomized controlled clinical trial)³⁵⁸

Outcome	Active events/ randomized	Control events/ randomized	Relative risk (95% CI)	Absolute risk difference (95% CI)	Number needed to treat (95% CI)	Attributable events (95% CI)
Very satisfied or satisfied with the treatment	16/17	16/18	1.06 (0.87; 1.30)	0.05 (-0.13; 0.24)		
Experienced side-effects	8/17	5/18	1.69 (0.69; 4.16)	0.19 (-0.12; 0.51)		
Constipation	1/17	0/18	3.17 (0.14; 72.80)	0.06 (-0.09; 0.21)		
Hematuria secondary to nephrolithiasis	1/17	0/18	3.17 (0.14; 72.80)	0.06 (-0.09; 0.21)		
Urinary tract infection	1/17	2/18	0.53 (0.05; 5.32)	-0.05 (-0.24; 0.13)		
Vaginal discomfort	0/17	2/18	0.21 (0.01; 4.10)	-0.11 (-0.28; 0.06)		
Vaginal hemorrhage	0/17	1/18	0.35 (0.02; 8.09)	-0.06 (-0.20; 0.09)		
Xerostomia	5/17	0/18	11.61 (0.69; 195.26)	0.29 (0.07; 0.52)	3 (14; 2)	294 (69; 519)