

Appendix Table F27. Pharmacological treatments for female UI (continued)

Reference study, country, sample	Inclusion criteria	Exclusion criteria	Active	Control	Sponsorship	Conflict of interest
Zinner, 2011 ⁴¹⁰ RCT Not reported N: 944	Male and female subjects aged ≥ 18 years with symptoms of OAB for ≥ 6 months who met the following criteria (based on a 3-day patient diary): urinary frequency ≥ 30 toilet voids/3 days (i.e. average ≥ 10 toilet voids/day); ≥ 1 "severe" urgency severity rating/3 days (as measured by the Indevus Urgency Severity Scale [IUSS]); and ≥ 3 UUI episodes/3 days (i.e., average ≥ 1 UUI episodes/day).	Subjects with a total voided volume >3000 ml/day or a mean volume voided/void >250 ml ; subjects with predominantly stress, insensate, or overflow incontinence; history of neurogenic bladder, indwelling or intermittent catheterization, significant renal disease (serum creatinine >1.5 mg /dL), uninvestigated hematuria or urinary tract infection during screening, or a history of ≥ 3 urinary tract infections in the previous 12 months; clinically significant urinary retention (defined as post-void residual urine volume >100 mL), cancer, interstitial cystitis.	Trospium for 48 weeks	Trospium for 36 weeks	Sponsored by Allergan Inc. and Endo Pharmaceuticals (formerly Indevus Pharmaceuticals, Inc.). Neil Reynolds, Monica Grandison, and Sushma Soni of in Science communications provided editorial support funded by Allergan, Inc.	None

Abbreviation: NR = Not reported