

Appendix Table F66. Quality of life after botulinum toxin vs. placebo, individual RCTs

Reference	Dose	Outcome	Active N	Control N	Active mean+/- standard deviation	Control mean+/- standard deviation	Mean difference	Lower 95% CI	Upper 95% CI
Ghei, 2005 ²⁹²	5000IU	KHQ score: emotional problems	10	10	5.3+/-2.02	7.0+/-2.42	-1.75	-3.70	0.20
Ghei, 2005 ²⁹²	5000IU	KHQ score: impact on life	10	10	1.5+/-0.81	2.5+/-0.81	-1	-1.71	-0.29
Ghei, 2005 ²⁹²	5000IU	KHQ score: incontinence impact	10	10	4.5+/-3.23	7.0+/-4.03	-2.5	-5.70	0.70
Ghei, 2005 ²⁹²	5000IU	KHQ score: incontinence severity measures	10	10	8.5+/-3.23	12.0+/-4.03	-3.5	-6.70	-0.30
Ghei, 2005 ²⁹²	5000IU	KHQ score: personal relationships	10	10	2.0+/-4.03	3.5+/-3.23	-1.5	-4.70	1.70
Ghei, 2005 ²⁹²	5000IU	KHQ score: physical/social limitations	10	10	5.0+/-2.42	7.5+/-4.03	-2.5	-5.42	0.42
Ghei, 2005 ²⁹²	5000IU	KHQ score: present health	10	10	1.0+/-0.81	1.5+/-0.81	-0.5	-1.21	0.21
Ghei, 2005 ²⁹²	5000IU	KHQ score: role limitations	10	10	2.5+/-1.61	3.5+/-1.61	-1	-2.41	0.41
Ghei, 2005 ²⁹²	5000IU	KHQ score: sleep/energy disturbances	10	10	3.5+/-1.61	5.0+/-0.81	-1.5	-2.62	-0.38