

**Appendix Table F75. Comparative effectiveness of drugs on continence**

Reference	Active drug	Dose	Control drug	Dose	Active n/N	Control n/N	Relative risk (95% CI)	Absolute risk difference (95% CI)
Kaplan, 2010 <sup>322</sup>	Fesoterodine	4 to 8mg once daily	Tolterodine ER	4 mg once daily	609/963	566/974	1.1(1.0; 1.2)	0.05 (0.01; 0.09)
NCT00444925 <sup>58</sup>	Fesoterodine	4 to 8mg once daily	Tolterodine ER	4 to 8mg once daily	396/685	358/690	1.1(1.0; 1.2)	0.06 (0.01; 0.11)
Milani, 1993 <sup>348</sup>	Flavoxate	1200	Oxybutynin	5mg t.i.d.	14/50	21/50	0.7(0.4; 1.2)	-0.14 (-0.32; 0.04)
Diokno, 2003 <sup>273</sup>	Oxybutynin	10mg/d	Tolterodine	4mg/d	90/391	67/399	1.4(1.0; 1.8)	0.06 (0.01; 0.12)
Chapple, 2005 <sup>61</sup>	Solifenacin	5-10mg od	Tolterodine	4mg once daily	341/578	294/599	1.2(1.1; 1.3)	0.10 (0.04; 0.16)
Halaska, 2003 <sup>302</sup>	Trospium	20mg twice daily	Oxybutynin	5mg twice daily	60/267	11/90	1.8(1.0; 3.3)	0.10 (0.02; 0.19)