## Data table for Figure 13. Vaccination coverage for combined series (4:3:1:3\*:3:1:4) among children aged 19–35 months, by selected characteristics: United States, 2010 and 2017

Excel and PowerPoint: https://www.cdc.gov/nchs/hus/contents2018.htm#Figure\_013

|  | 2010    |                | 2017    |                |
|--|---------|----------------|---------|----------------|
|  | Percent | Standard error | Percent | Standard error |
| Total  | 56.6    | 0.7            | 70.4    | 0.8            |
| Race and Hispanic origin                       |         |                |         |                |
| White, not Hispanic                            | 56.9    | 0.8            | 71.5    | 0.9            |
| Black, not Hispanic                            | 54.5    | 1.8            | 66.5    | 2.2            |
| Asian, not Hispanic                            | 59.3    | 3.7            | 72.4    | 3.4            |
| American Indian or Alaska Native, not Hispanic | 64.1    | 4.7            | 65.9    | 5.1            |
| Hispanic                                       | 55.5    | 1.6            | 70.4    | 1.8            |
| Metropolitan statistical area (MSA)            |         |                |         |                |
| Non-MSA  | 55.2    | 1.5            | 66.8    | 1.6            |
| MSA, nonprincipal city                         | 57.7    | 1.1            | 69.8    | 1.2            |
| MSA, principal city                            | 56.1    | 1.1            | 71.9    | 1.1            |
| Health insurance coverage                      |         |                |         |                |
| Uninsured                                      | 41.0    | 3.6            | 48.5    | 3.7            |
| Medicaid                                       | 53.6    | 1.1            | 66.5    | 1.2            |
| Private  | 61.0    | 0.8            | 76.0    | 1.0            |

NOTES: Data are for the civilian noninstitutionalized population. The combined 7-vaccine series consists of 4 or more doses of either the diphtheria, tetanus toxoids, and pertussis vaccine (DTP), the diphtheria and tetanus toxoids vaccine (DT), or the diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP); 3 or more doses of any poliovirus vaccine; 1 or more doses of a measles-containing vaccine (MCV); 3 or more doses or 4 or more doses of *Haemophilus influenzae* type b vaccine (Hib) depending on Hib vaccine product type (full series Hib); 3 or more doses of hepatitis B vaccine; 1 or more doses of varicella vaccine; and 4 or more doses of pneumococcal conjugate vaccine (PCV). Persons of Hispanic origin may be of any race. Starting with 2000 data, estimates were tabulated using the 1997 Revisions to the *Standards for the Classification of Federal Data on Race and Ethnicity*. MSA status was grouped into three categories: MSA principal city, MSA non-principal city, and non-MSA. MSA and principal city were as defined by the U.S. Census Bureau (https://www.census.gov/geo/reference/gtc/gtc\_cbsa.html). Data from U.S. territories were not included in the national estimates. See Appendix II, Vaccination.

SOURCE: National Center for Immunization and Respiratory Diseases (NCIRD), National Immunization Survey-Child (NIS-Child). See Appendix I, National Immunization Surveys (NIS).

Health, United States, 2018