

Data table for Figure 2. Infant mortality rates, by race and Hispanic origin of mother and leading causes of death: United States, 2007–2017

Excel and PowerPoint: https://www.cdc.gov/nchs/hus/contents2018.htm#Figure_002

Infant mortality rates, by race and Hispanic origin, 2007–2017

Year	Race and Hispanic origin ¹					
	All mothers	Hispanic or Latina	Not Hispanic or Latina			
			White	Black or African American	Asian or Pacific Islander	American Indian or Alaska Native
Infant deaths per 1,000 live births ²						
2007	6.75	5.51	5.63	13.31	4.60	9.37
2008	6.61	5.59	5.53	12.67	4.39	8.67
2009	6.39	5.29	5.33	12.40	4.27	9.17
2010	6.14	5.25	5.18	11.46	4.18	8.64
2011	6.07	5.15	5.07	11.45	4.18	8.52
2012	5.98	5.11	5.04	11.19	3.96	8.73
2013	5.96	5.00	5.06	11.11	3.90	7.72
2014	5.82	5.01	4.89	10.93	3.68	7.66
2015	5.90	4.96	4.90	11.25	4.08	8.58
2016	5.87	4.96	4.87	11.21	3.83	8.81
2017	5.79	5.10	4.69	10.88	4.03	8.90
Standard error						
2007	0.04	0.07	0.05	0.14	0.14	0.47
2008	0.04	0.07	0.05	0.15	0.14	0.45
2009	0.04	0.07	0.05	0.14	0.13	0.47
2010	0.04	0.07	0.05	0.14	0.13	0.47
2011	0.04	0.08	0.05	0.14	0.13	0.47
2012	0.04	0.08	0.05	0.14	0.12	0.48
2013	0.04	0.07	0.05	0.14	0.12	0.45
2014	0.04	0.07	0.05	0.14	0.12	0.45
2015	0.04	0.07	0.05	0.14	0.12	0.48
2016	0.04	0.07	0.05	0.14	0.12	0.50
2017	0.04	0.08	0.05	0.14	0.12	0.51

See footnotes at end of table.

Data table for Figure 2. Infant mortality rates, by race and Hispanic origin of mother and leading causes of death: United States, 2007–2017—Con.

Excel and PowerPoint: https://www.cdc.gov/nchs/hus/contents2018.htm#Figure_002

Infant mortality rates, by leading causes of death: 2007–2017

Year	Leading causes of death ³				
	Congenital malformations	Preterm births and low birthweight	Sudden infant death syndrome	Maternal complications of pregnancy	Unintentional injuries (accidents)
	Infant deaths per 100,000 live births ²				
2007	134.9	112.7	57.0	41.1	29.7
2008	133.7	112.0	55.3	41.8	30.9
2009	129.7	109.6	54.0	39.1	28.4
2010	127.9	103.8	51.5	39.1	27.7
2011	126.9	104.1	48.2	40.4	29.5
2012	125.7	106.6	42.4	38.4	29.4
2013	121.5	107.1	39.7	40.6	29.3
2014	119.2	104.6	38.6	39.6	29.2
2015	121.8	102.7	39.4	38.4	32.4
2016	122.2	99.5	38.0	35.7	30.8
2017	119.2	97.5	35.3	37.2	34.1
	Standard error				
2007	1.77	1.62	1.15	0.98	0.83
2008	1.78	1.62	1.14	0.99	0.85
2009	1.77	1.63	1.14	0.97	0.83
2010	1.79	1.61	1.13	0.99	0.83
2011	1.79	1.62	1.10	1.01	0.86
2012	1.78	1.64	1.04	0.99	0.86
2013	1.76	1.65	1.00	1.02	0.86
2014	1.73	1.62	0.98	1.00	0.86
2015	1.75	1.61	1.00	0.98	0.90
2016	1.76	1.59	0.98	0.95	0.88
2017	1.76	1.59	0.96	0.98	0.94

¹Starting with 2003 data, some states reported multiple-race data for births and deaths according to the 1997 OMB standards. The multiple-race data for these states were bridged to the four single-race categories of the 1977 OMB standards, for comparability across the trend. Persons of Hispanic origin may be of any race. See [Appendix I](#), Population Census and Population Estimates; [Appendix II](#), Hispanic origin; Race.

²Infants are defined as under 1 year of age. Rates are based on the number of deaths from the mortality file and the number of births from the natality file.

³Deaths are identified using *International Classification of Diseases, 10th revision (ICD–10)*. The codes were: Congenital malformations, deformations and chromosomal abnormalities (Q00–Q99); Disorders related to preterm births (short gestation) and low birthweight, not elsewhere classified (P07); Sudden infant death syndrome (R95); Newborn affected by maternal complications of pregnancy of pregnancy (P01); and Accidents (unintentional injuries) (V01–X59). The cause of death, newborn affected by maternal complications of pregnancy, includes any of the following conditions: incompetent cervix, premature rupture of membranes, ectopic pregnancy, malpresentation before labor, amniotic fluid disorders, multiple pregnancy, maternal death, and other unspecified maternal complications. See [Appendix II](#), Cause of death; Table IV.

NOTES: Infant mortality rates by race and Hispanic origin are from the Linked Birth/Infant Death data set, for which the most recent data year available is 2017. Age-adjusted rates are calculated using the year 2000 standard population with unrounded population numbers. See [Appendix II](#), Cause of death; Table IV; Hispanic origin; Race.

SOURCE: NCHS, National Vital Statistics System, Linked Birth/Infant Death Data Set. See [Appendix I](#), National Vital Statistics System (NVSS).