



## Mebendazole

Updated: June 6, 2019.

## OVERVIEW

### Introduction

Mebendazole is an antihelmintic agent used commonly for roundworm (pinworm and hookworm) infections, trichinosis, capillariasis and toxocariasis and other parasitic worm infections. Mebendazole when given for prolonged periods in high doses has been associated with elevations in serum enzyme levels, and rare instances of acute, clinically apparent liver injury have been linked to its use.

### Background

Mebendazole (me ben' da zole) is a benzimidazole antihelmintic similar in structure and mechanism of action to thiabendazole and albendazole. The benzimidazoles act by selective binding to beta-tubulin of parasitic worms, causing their immobilization and death. Mebendazole and albendazole share similar antihelmintic activity, mebendazole generally being preferred for treatment of pinworm and roundworm infections, and albendazole (because it is better absorbed) for systemic parasitic infections such as echinococcosis and cysticercosis. Mebendazole was approved for use in the United States in 1974 and is indicated for therapy of common parasitic worm infections. Mebendazole is available generically in 100 mg chewable tablets. The usual dose is 100 to 500 mg once (pinworm) or varying doses for 3 days (whipworm, hookworm and roundworm infections), or varying doses for up to 28 days, depending upon the indication. Side effects are uncommon, but can include gastrointestinal upset, fever and diarrhea.

### Hepatotoxicity

Mebendazole when given in typical doses has not been associated with serum enzyme elevations, although the duration of therapy is usually short and monitoring for enzyme elevations has rarely been reported. With high dose therapy (which is now rarely used with the availability of albendazole), elevations in serum aminotransferase levels (2 to 10 times normal) can occur, but are usually well tolerated. There have been rare reports of acute liver injury due to mebendazole, particularly when it is given repeatedly or in higher doses. The onset is usually with fever and malaise within days of starting or restarting therapy. The pattern of serum enzyme elevations is typically hepatocellular and jaundice is uncommon. The abnormalities usually resolve rapidly with stopping therapy. Signs of hypersensitivity (rash, fever and eosinophilia) are typical and liver biopsy may show granulomas.

Likelihood score: D (possible cause of clinically apparent liver injury).

## Mechanism of Injury

Mebendazole acts by binding tubulin in parasitic worms, which it does with greater avidity than the tubulin in mammalian cells, but some of the toxicity of the benzimidazoles may be related to this tubulin-binding activity. In most instances of clinically apparent liver injury, hypersensitivity appears to be the cause.

## Outcome and Management

Mebendazole is usually well tolerated and the liver injury reported with its use has been mild and self-limited in course. Patients with hypersensitivity and acute liver injury attributed to mebendazole should avoid repeat exposure. It is unknown whether there is cross sensitivity with other benzimidazoles (such as albendazole), but there probably is and switching to another class of antihelmintic agents is appropriate if therapy is still needed.

Drug Class: [Antihelmintic Agents](#)

## CASE REPORT

### Case 1. Acute liver injury due to mebendazole.

[Modified from: Colle I, Naegels S, Hoorens A, Hautekeete M. Granulomatous hepatitis due to mebendazole. *J Clin Gastroenterol* 1999; 28: 44-5. [PubMed Citation](#)]

A 52 year old Belgian man was treated with mebendazole (100 mg twice daily) for 3 days for suspected ascariasis. Fourteen days later the course was repeated, but within 2 days of restarting mebendazole he developed fever (39 °C), diarrhea, poor appetite and fatigue. Because of the fever, he was given a 5 day course of cefuroxime, but remained febrile and symptomatic and was then found to have elevations in serum aminotransferase levels, with normal serum bilirubin and alkaline phosphatase (Table). He had eosinophilia (18%: absolute count 2,286/ $\mu$ L). Tests for hepatitis A, B and C were negative. He had low levels of antinuclear antibody (ANA: 1:40) and smooth muscle antibody (SMA: 1:160), but total globulin levels were normal. Ultrasound of the abdomen was negative. Stools were negative for ova and parasites. A liver biopsy showed multiple granulomas with multinucleate cells and active inflammation. There were no ova or helminths visualized and special stains for mycobacteria were negative. He improved rapidly without further therapy and 3 months later serum enzymes were normal.

### Key Points

Medication:	Mebendazole (100 mg twice daily) for ~5 days
Pattern:	Hepatocellular (R=9.7)
Severity:	1+ (enzyme elevations without jaundice)
Latency:	26 days after initial course, 2 days of second course
Recovery:	Within 3 months
Other medications:	Cefuroxime (after onset of symptoms)

### Laboratory Values

Days After Starting	Days After Stopping	ALT (U/L)	Alk P (U/L)	Bilirubin (mg/dL)	Other
0	Mebendazole (100 mg twice daily for 3 days)				
14	Mebendazole (100 mg twice daily for 2 days)				
16	0	Fever and rash			

Table continued from previous page.

Days After Starting	Days After Stopping	ALT (U/L)	Alk P (U/L)	Bilirubin (mg/dL)	Other
26	10	466	170	1.0	Eosinophilia
28	12	540			Liver biopsy
100	90	Normal	Normal	Normal	No symptoms
<b>Normal Values</b>		<b>&lt;48</b>	<b>&lt;60</b>	<b>&lt;1.2</b>	

## Comment

Acute hepatocellular injury arose concurrent with symptoms and signs of hypersensitivity (rash and fever) within days of starting a second course of mebendazole. This pattern is typical of immunoallergic hepatitis and usually resolves rapidly, but can be severe if there is reexposure. The ANA and SMA titers may have been stimulated by the acute injury; in the absence of hyperglobulinemia were not compatible with autoimmune hepatitis. The granulomas found on liver biopsy reflect the generalized hypersensitivity and similar granulomas are likely to be found in other organs (lymph nodes, spleen).

## PRODUCT INFORMATION

### REPRESENTATIVE TRADE NAMES

Mebendazole – Generic


### DRUG CLASS

Antihelminthic Agents

### COMPLETE LABELING

Product labeling at DailyMed, National Library of Medicine, NIH

## CHEMICAL FORMULA AND STRUCTURE

DRUG	CAS REGISTRY NUMBER	MOLECULAR FORMULA	STRUCTURE
Mebendazole	31431-39-7	C <sub>16</sub> -H <sub>13</sub> -N <sub>3</sub> -O <sub>3</sub>	

## ANNOTATED BIBLIOGRAPHY

References updated: 6 June 2019

Zimmerman HJ. Antihelminthics. Hepatic injury from antimicrobial agents. In, Zimmerman HJ. Hepatotoxicity: the adverse effects of drugs and other chemicals on the liver. 2nd ed. Philadelphia: Lippincott, 1999, pp. 626-8.

- (Expert review of hepatotoxicity of antihelmintics written in 1999; mebendazole produces hepatic injury in dogs and occasionally in humans by uncertain mechanisms).*
- Keiser J, McCarthy J, Hotez PJ. Chemotherapy of helminth infections. In, Brunton LL, Hilal-Dandan R, Knollman BC, eds. Goodman & Gilman's the pharmacological basis of therapeutics. 13th ed. New York: McGraw-Hill, 2018, pp. 1001-10.
- (Textbook of pharmacology and therapeutics).*
- Polzin DJ, Stowe CM, O'Leary TP, Stevens JB, Hardy RM. Acute hepatic necrosis associated with the administration of mebendazole to dogs. J Am Vet Med Assoc 1981; 179: 1013-6. PubMed PMID: 7341557.
- (Four of 10 Dachshunds treated with mebendazole [22 mg/kg for 3-5 days] developed acute hepatitis 8-10 days after stopping therapy [bilirubin 14.2 mg/dL, ALT 17 times ULN, Alk P 8 times ULN], and two died of multiorgan failure, autopsy showing centrilobular coagulative necrosis).*
- Swanson JE, Breider MA. Hepatic failure following mebendazole administration to a dog. J Am Vet Med Assoc 1982; 181: 72-3. PubMed PMID: 7107495.
- (4 year old Labrador Retriever developed liver injury [bilirubin 8.1 mg/dL, ALT 8 times ULN, Alk P 5 times ULN] starting 3 days after a second course of mebendazole [22 mg/kg], autopsy showing centrilobular hepatic necrosis).*
- Seitz R, Schwerk W, Arnold R. [Hepatocellular drug reaction caused by mebendazole therapy in cystic echinococcosis]. Z Gastroenterol 1983; 21: 324-9. German. PubMed PMID: 6613213.
- (76 year old man with Echinococcal cysts of liver developed rash 49 days after starting mebendazole [~29 mg/kg/day], with ALT 250 U/L but normal bilirubin and Alk P; positive rechallenge to low dose [ALT 403 U/L]).*
- Junge U, Mohr W. [Mebendazole-hepatitis]. Z Gastroenterol 1983; 21: 736-8. German. PubMed PMID: 6666188.
- (Among 29 patients with echinococcosis treated with mebendazole, 1 developed hepatitis; 53 year old woman noted dark urine 2 months after starting therapy [bilirubin 2.1 mg/dL, ALT 475 U/L, Alk P 270 U/L]; biopsy showing spotty necrosis, resolving in 1.5 months; positive rechallenge [ALT 400 U/L, Alk P 358 U/L]).*
- Mousa AM, Rudwan MA, Marafi AA, Muhtaseb SA, Dajani AI. Human cystic hydatid disease: treatment with interrupted courses of mebendazole. J Trop Med Hyg 1986; 89: 257-64. PubMed PMID: 3795326.
- (10 patients with echinococcal cysts were treated with high doses of mebendazole [1800 mg/day] for up to 18 weeks; a 37 year old woman with multiple cysts developed jaundice at 11 days that became cholestatic but ultimately resolved; few details given and referred to as "amoebic hepatitis").*
- Bekhti A, Pirotte J. Hepatotoxicity of mebendazole. Relationship with serum concentrations of the drug. Gastroenterol Clin Biol 1987; 11: 701-3. PubMed PMID: 3692093.
- (47 year old woman with hydatid cysts in the liver developed persistent elevations in ALT [60-400 U/L] and Alk P [207 U/L] without jaundice during a third course of mebendazole given in high doses [4.5-6.0 g daily], ALT levels correlated with serum mebendazole levels, normalized with stopping and rose again with rechallenge apparently without symptoms or jaundice).*
- Colle I, Naegels S, Hoorens A, Hautekeete M. Granulomatous hepatitis due to mebendazole. J Clin Gastroenterol 1999; 28: 44-5. PubMed PMID: 916665.
- (52 year old man treated with mebendazole for 3 days on two occasions 2 weeks apart developed fever during the second course and then found to have bilirubin 1.0 mg/dL, ALT 466 U/L, Alk P normal, 18% eosinophils and liver biopsy showing granulomas; no jaundice and rapid recovery: Case 1).*

Reuter S, Jensen B, Buttenschoen K, Kratzer W, Kern P. Benzimidazoles in the treatment of alveolar echinococcosis: a comparative study and review of the literature. *J Antimicrob Chemother* 2000; 46: 451-6. PubMed PMID: 10980173.

*(35 patients with alveolar echinococcosis were treated with either mebendazole or albendazole for 12-79 months; no severe side effects, but one patient was switched from albendazole to mebendazole because of ALT elevations).*

Keiser J, Utzinger J. Efficacy of current drugs against soil-transmitted helminth infections: systematic review and meta-analysis. *JAMA* 2008; 299: 1937-48. PubMed PMID: 18430913.

*(Systematic review of efficacy and safety of albendazole, mebendazole and pyrantel pamoate as therapy of *Ascaris lumbricoides* [roundworm], hookworm and whipworm; mebendazole was well tolerated in most studies, no significant adverse events were reported).*

Drugs for parasitic infections. 2nd ed. New Rochelle, NY: Medical Letter, Inc., 2010.

*(Thorough description of drugs for parasitic infections in adults and children as well as a table of their major side effects; mebendazole can be used for ascariasis, pinworm, hookworm, whipworm, trichinellosis, and several less common parasitic infections; no hepatic related side effects mentioned).*

Mandal S, Mandal MD. Human cystic echinococcosis: epidemiologic, zoonotic, clinical, diagnostic and therapeutic aspects. *Asian Pac J Trop Med* 2012; 5: 253-60. PubMed PMID: 22449514.

*(Review of human echinococcal cyst disease, its epidemiology, clinical manifestations, diagnosis, prevention and treatment which includes direct aspiration and surgery on the cysts and medical therapy with either mebendazole, albendazole or ivermectin often with praziquantel; no discussion of side effects).*

Chalasani N, Bonkovsky HL, Fontana R, Lee W, Stolz A, Talwalkar J, Reddy KR, et al.; United States Drug Induced Liver Injury Network. Features and outcomes of 899 patients with drug-induced liver injury: The DILIN Prospective Study. *Gastroenterology* 2015; 148: 1340-52. PubMed PMID: 25754159.

*(Among 899 cases of drug induced liver injury enrolled in a US prospective study between 2004 and 2013, no cases were attributed to mebendazole or other antihelminthic medications).*