



Nelarabine

Updated: July 10, 2014.

OVERVIEW

Introduction

Nelarabine is a purine analogue and antineoplastic agent used in the therapy of T cell lymphoblastic leukemia or lymphoma. Nelarabine is associated with a low rate of transient serum enzyme elevations during therapy and has been linked to rare instances of clinically apparent acute liver injury with jaundice.

Background

Nelarabine (ne lar' a been) is a purine analogue that is used in the treatment of T cell leukemia or lymphoma. Nelarabine is arabinosyl derivative of deoxyguanosine (2-amino-9-beta-D-arabinofuranosyl-6-methoxy-9-H-purine) that, after removal of the methoxy group by adenosine deaminase in serum, is taken up and converted intracellularly to the active triphosphate, which is believed to compete with guanine triphosphate for use by DNA polymerase leading to inhibition of DNA synthesis. It has selective activity against T lymphocytes and was found to have activity against acute T cell malignancies. Nelarabine was approved for use as an antineoplastic agent in the United States in 2005. Current indications are therapy of acute T cell lymphoblastic leukemia and T cell lymphoblastic lymphoma after failure of prior therapies. Nelarabine is available as a solution for injection under the trade name Arranon. The typical adult dose is 1500 mg/m² intravenously on days 1, 3 and 5 of 21-day cycles. Common side effects include bone marrow suppression, nausea, vomiting, anorexia, diarrhea, headache, fatigue, mucositis and skin rash. The major dose limiting side effects of nelarabine are neurologic, including somnolence, headache, dizziness, ataxia, delirium, seizures, neuropathy and Guillain Barre syndrome. The neurologic toxicities can be severe and some are not reversible on stopping nelarabine.

Hepatotoxicity

In clinical trials, serum enzymes elevations occurred in a small proportion of patients treated with nelarabine when given as sole therapy for refractory or relapsed acute leukemia. These elevations are generally mild-to-moderate, transient and asymptomatic. Elevations of aminotransferase levels above 5 times the upper limit of normal are reported in 4% of patients with leukemia receiving nelarabine. The elevations rarely require dose adjustment or delay in therapy. Cases of clinically apparent liver injury due to nelarabine have been reported to occur, but few details are available. A single case report of clinically apparent liver injury attributed to nelarabine has been published with rapid onset of jaundice during a second course of nelarabine, a hepatocellular pattern of enzyme elevations, no immunoallergic or autoimmune features and a rapid improvement upon stopping.

Mechanism of Injury

Hepatotoxicity from nelarabine is likely due to direct toxicity as is typical for other purine analogues.

Outcome and Management

The severity of the liver injury linked to nelarabine therapy is generally self-limited and mild and resolves with stopping therapy. There is little evidence of cross sensitivity to liver injury among the various antineoplastic or antiviral purine analogues.

Drug Class: [Antineoplastic Agents](#), Antimetabolites

Other Drugs in the Subclass, [Purine Analogues](#): Azathioprine, Cladribine, Clofarabine, Fludarabine, Mercaptopurine, Pentostatin, Thioguanine

PRODUCT INFORMATION

REPRESENTATIVE TRADE NAMES

Nelarabine – Generic, Arranon®


DRUG CLASS

Antineoplastic Agents

COMPLETE LABELING

Product labeling at DailyMed, National Library of Medicine, NIH

CHEMICAL FORMULA AND STRUCTURE

DRUG	CAS REGISTRY NUMBER	MOLECULAR FORMULA	STRUCTURE
Nelarabine	121032-29-9	C ₁₁ -H ₁₅ -N ₅ -O ₅	

ANNOTATED BIBLIOGRAPHY

References updated: 10 July 2014

Zimmerman HJ. Oncotherapeutic and immunosuppressive agents. In, Zimmerman HJ. Hepatotoxicity: the adverse effects of drugs and other chemicals on the liver. 2nd ed. Philadelphia: Lippincott, 1999, pp. 673-708.

(Expert review of hepatotoxicity of cancer chemotherapeutic agents published in 1999 discusses the purine analogues pentostatin, cladribine and fludarabine but not clofarabine or nelarabine).

DeLeve LD. Cancer chemotherapy. In, Kaplowitz N, DeLeve LD, eds. Drug-induced liver disease. 3rd ed. Amsterdam: Elsevier, 2013, p. 541-68.

(Review of hepatotoxicity of anticancer agents does not discuss nelarabine).

Chabner BA, Bertino J, Cleary J, Ortiz T, Lane A, Supko JG, Ryan DP. Purine analogs. Cytotoxic agents. Chemotherapy of neoplastic diseases. In, Brunton LL, Chabner BA, Knollman BC, eds. Goodman & Gilman's the pharmacological basis of therapeutics. 12th ed. New York: McGraw-Hill, 2011, pp. 1701-5.

(Textbook of pharmacology and therapeutics).

Kisor DF. Nelarabine: a nucleoside analog with efficacy in T-cell and other leukemias. Ann Pharmacother 2005; 39: 1056-63. PubMed PMID: 15870141.

(Analysis of structure, mechanism of action, pharmacokinetics, efficacy and safety of nelarabine in T cell malignancies mentions that the dose limiting toxicity is neurological; no discussion of hepatotoxicity).

Gandhi V, Plunkett W. Clofarabine and nelarabine: two new purine nucleoside analogs. Curr Opin Oncol 2006; 18: 584-90. PubMed PMID: 16988579.

(Review of clofarabine and nelarabine, both of which had been recently approved for use in ALL and T cell leukemia or lymphoma, dose limiting toxicities being hepatic for clofarabine, neurologic for nelarabine).

Czuczman MS, Porcu P, Johnson J, Niedzwiecki D, Kelly M, Hsi ED, Cook JR, et al.; Cancer and Leukemia Group B. Results of a phase II study of 506U78 in cutaneous T-cell lymphoma and peripheral T-cell lymphoma: CALGB 59901. Leuk Lymphoma 2007; 48: 97-103. PubMed PMID: 17325852.

(Among 18 adults with T cell lymphomas treated with nelarabine, none had ALT or AST elevations during therapy, but neurologic toxicity was common [33%] and dose limiting).

Buie LW, Epstein SS, Lindley CM. Nelarabine: a novel purine antimetabolite antineoplastic agent. Clin Ther 2007; 29: 1887-99. PubMed PMID: 18035189.

(Review of the structure, mechanism of action, pharmacokinetics, clinical efficacy and safety of nelarabine encountered in preapproval trials; discusses the neurologic and hematologic, but not hepatic toxicities of nelarabine).

Cohen MH, Johnson JR, Justice R, Pazdur R. FDA drug approval summary: nelarabine (Arranon) for the treatment of T-cell lymphoblastic leukemia/lymphoma. Oncologist 2008; 13: 709-14. PubMed PMID: 18586926.

(Summary of the evidence for efficacy and safety of nelarabine in T cell leukemia and lymphoma which led to its approval in refractory or relapsed cases; the major dose limiting toxicity was neurologic, elevations of ALT levels above 5 times the ULN occurred in 4% of patients; no mention of clinically apparent liver injury).

Iino M. [Severe liver injury following nelarabine chemotherapy for T-cell lymphoblastic lymphoma]. Rinsho Ketsueki 2009; 50: 49-51. Japanese. PubMed PMID: 19225230.

(41 year old man with T cell lymphoma after HCT developed liver test abnormalities at the end of a second course of nelarabine [peak bilirubin 17.7 mg/dL, ALT 2149 U/L, Alk P 704 U/L], resolving within 1 month of stopping with prednisolone and ursodiol treatment).

Commander LA, Seif AE, Insogna IG, Rheingold SR. Salvage therapy with nelarabine, etoposide, and cyclophosphamide in relapsed/refractory paediatric T-cell lymphoblastic leukaemia and lymphoma. *Br J Haematol* 2010; 150: 345-51. PubMed PMID: 20528871.

(Among 7 children with T cell leukemia treated with nelarabine in combination with etoposide and cyclophosphamide, 6 developed neurological toxicities, mostly pain, headache, somnolence, dizziness, neuropathy, tremor and ataxia; no mention of ALT elevations or hepatotoxicity).

Dunsmore KP, Devidas M, Linda SB, Borowitz MJ, Winick N, Hunger SP, Carroll WL, et al. Pilot study of nelarabine in combination with intensive chemotherapy in high-risk T-cell acute lymphoblastic leukemia: a report from the Children's Oncology Group. *J Clin Oncol* 2012; 30: 2753-9. PubMed PMID: 22734022.

(Among 92 children with T cell leukemia, 72 received nelarabine in addition of standard therapy; ALT and AST elevations occurred in 44% on nelarabine and in 56% who were not; neurologic side effects were more common with nelarabine, 11 developing neuropathy, 4 seizures and 1 a Guillain Barre-like syndrome).