



Pentostatin

Updated: July 10, 2014.

OVERVIEW

Introduction

Pentostatin is a purine analogue and antineoplastic agent used in the therapy of hairy cell leukemia and T cell lymphomas. Pentostatin is associated with a low rate of serum enzyme elevations during therapy and has been linked to rare instances of severe acute liver injury with jaundice.

Background

Pentostatin (pen" toe stat' in) is a purine analogue (2'-deoxycoformycin) that is used in the treatment of hairy cell leukemia and T cell lymphomas. Pentostatin is a transition state analogue of a major intermediate in the pathway of the adenosine deaminase reaction. As a consequence, pentostatin inhibits adenosine deaminase and causes accumulation of intracellular adenosine and deoxyadenosine which cause a block DNA synthesis. This imbalance of nucleotide pools is particularly toxic to lymphocytes. Pentostatin was found to have marked activity against hairy leukemia and T cell lymphomas and was approved for this use in the United States in 1991. Pentostatin is available as a solution for injection generically and under the trade name Nipent. The typical dose regimen is 4 mg/m² given intravenously every other week for 6 months. Repeat courses are recommended only for patients who have a response and later relapse. Common side effects include bone marrow suppression, fever, infections, nausea, vomiting, anorexia, diarrhea, headache, fatigue and skin rash. Because pentostatin tends to deplete normal T cells, opportunistic infections can occur and immunosuppression can persist for months or years after discontinuation. At high doses, pentostatin can cause severe renal and neurologic toxicity and in combination with fludarabine can cause pulmonary toxicity.

Hepatotoxicity

In clinical trials, serum enzymes elevations occurred in up to 25% of patients receiving pentostatin, but the abnormalities were generally mild and transient and rarely required dose modification. Clinically apparent liver injury from pentostatin is rare, but striking instances of severe liver injury leading rapidly to multiorgan failure and death have been described both in adults and children. The time to onset varied from a few days to six months. The possible role of shock, ischemia, opportunistic infections and sepsis in these cases has not always been well defined. Both hepatocellular and cholestatic patterns of enzyme elevations have been described. Immunoallergic features and autoantibodies were not typical.

Mechanism of Injury

In several instances, hepatic toxicity from pentostatin appeared to be somewhat dose related, suggesting that the liver injury is a direct effect of the purine analogue. Because pentostatin is a potent immunosuppressive agent,

the possibility exists that some cases of hepatic injury are due to reactivation of hepatitis B or other opportunistic infections. While pentostatin has not been shown to cause reactivation of hepatitis B, there is a strong possibility that it might induce this syndrome, and several cases of hepatic injury during pentostatin therapy were described as due to concurrent hepatitis B.

Drug Class: [Antineoplastic Agents](#), Antimetabolites

Other Drugs in the Subclass, [Purine Analogues](#): Azathioprine, Cladribine, Clofarabine, Fludarabine, Mercaptopurine, Nelarabine, Thioguanine

PRODUCT INFORMATION

REPRESENTATIVE TRADE NAMES

Pentostatin – Generic, Nipent®


DRUG CLASS

Antineoplastic Agents

COMPLETE LABELING

Product labeling at [DailyMed](#), National Library of Medicine, NIH

CHEMICAL FORMULA AND STRUCTURE

DRUG	CAS REGISTRY NUMBER	MOLECULAR FORMULA	STRUCTURE
Pentostatin	53910-25-1	C ₁₁ -H ₁₆ -N ₄ -O ₄	 <p>The chemical structure of Pentostatin is a complex molecule consisting of a 7-membered diazepine ring fused to a 5-membered imidazole ring. The diazepine ring has a secondary amine group (HN) and a nitrogen atom at the 2-position. The imidazole ring has a nitrogen atom at the 4-position. A hydroxyl group (OH) is attached to the 5-position of the diazepine ring. The 2-position of the imidazole ring is connected to a ribose sugar ring. The ribose sugar has hydroxyl groups (HO) at the 2' and 3' positions.</p>

ANNOTATED BIBLIOGRAPHY

References updated: 10 July 2014

Zimmerman HJ. Oncotherapeutic and immunosuppressive agents. In, Zimmerman HJ. Hepatotoxicity: the adverse effects of drugs and other chemicals on the liver. 2nd ed. Philadelphia: Lippincott, 1999, pp. 673-708.

(Expert review of hepatotoxicity of cancer chemotherapeutic agents published in 1999 mentions that pentostatin is an adenosine deaminase inhibitor, has hepatotoxic potential and causes aminotransferase elevations in both experimental animals and humans).

DeLeve LD. Cancer chemotherapy. In, Kaplowitz N, DeLeve LD, eds. Drug-induced liver disease. 3rd ed. Amsterdam: Elsevier, 2013, p. 541-68.

(Review of hepatotoxicity of anticancer agents does not discuss pentostatin).

Chabner BA, Bertino J, Cleary J, Ortiz T, Lane A, Supko JG, Ryan DP. Purine analogs. Cytotoxic agents. Chemotherapy of neoplastic diseases. In, Brunton LL, Chabner BA, Knollman BC, eds. Goodman & Gilman's the pharmacological basis of therapeutics. 12th ed. New York: McGraw-Hill, 2011, pp. 1701-5.

(Textbook of pharmacology and therapeutics).

Poplack DG, Sallan SE, Rivera G, Holcenberg J, Murphy SB, Blatt J, Lipton JM et al. Phase I study of 2'-deoxycoformycin in acute lymphoblastic leukemia. *Cancer Res* 1981; 41): 3343-6. PubMed PMID: 6973390.

(Among 26 children with ALL treated with 43 courses of pentostatin [0.25-1.0 mg/kg/day for 3 days], dose-limiting toxicities were neurologic [seizures, coma], pulmonary, renal and hepatic, including 2 patients receiving high doses who developed severe injury: one had jaundice and fever the day after completing the first course [bilirubin not given, ALT 743 U/L] and died of respiratory failure one day later; a second developed jaundice, fever and ascites after 2 days of therapy [bilirubin 15.4 mg/dL, ALT normal] and died two weeks later; autopsy in both patients failed to demonstrate a clear cause).

Spiers AS, Parekh SJ, Bishop MB. Hairy-cell leukemia: induction of complete remission with pentostatin (2'-deoxycoformycin). *J Clin Oncol* 1984; 2: 1336-42. PubMed PMID: 6334721.

(Two men, ages 46 and 61 years, with hairy cell leukemia were treated with pentostatin [7 to 8 courses of 5 mg/kg/day for 2 days] and both had a long lasting complete remission; skin rash, but no other toxicities were mentioned).

Dang-Vu AP, Olsen EA, Vollmer RT, Greenberg ML, Hershfield MS. Treatment of cutaneous T cell lymphoma with 2'-deoxycoformycin(pentostatin). *J Am Acad Dermatol* 1988; 19: 692-8. PubMed PMID: 3263401.

(3 patients with cutaneous T cell lymphoma were treated with pentostatin [5 mg/m²/day], one of whom developed side effects of episcleritis, nausea and fever within 4 days of starting with subsequent liver enzyme elevations [ALT rising from 9 to 38 U/L, Alk P from 38 to 120 U/L] that persisted for at least 10 months after stopping).

Ho AD, Thaler J, Stryckmans P, Coiffier B, Luciani M, Sonneveld P, Lechner K, et al. Pentostatin in refractory chronic lymphocytic leukemia: a phase II trial of the European Organization for Research and Treatment of Cancer. *J Natl Cancer Inst* 1990; 82: 1416-20. PubMed PMID: 2388293.

(Among 26 adults with refractory CLL treated with pentostatin [4 mg/m² every 1-2 weeks], liver biochemical tests became abnormal in 20%, but values over 5 times ULN occurred in only 1 patient who was diagnosed as having hepatitis B, but no details provided).

Monfardini S, Sorio R, Cavalli F, Cerny TH, Van Glabbeke M, Kaye S, Smyth JF. Pentostatin (2'-deoxycoformycin, dCF) in patients with low-grade (B-T-cell) and intermediate- and high-grade (T-cell)

malignant lymphomas: phase II study of the EORTC Early Clinical Trials Group. *Oncology* 1996; 53: 163-8. PubMed PMID: 8604244.

(Among 37 patients with low grade non-Hodgkin lymphoma treated with weekly or every other week 1-day courses of pentostatin [5 mg/m²/day], toxicities included leucopenia, infection, creatinine elevations, nausea, vomiting, diarrhea, conjunctivitis and rash).

Kurzrock R, Strom SS, Estey E, O'Brien S, Keating MJ, Jiang H, Adams T, et al. Second cancer risk in hairy cell leukemia: analysis of 350 patients. *J Clin Oncol* 1997; 15: 1803-10. PubMed PMID: 9164188.

(Among 350 patients with hairy cell leukemia treated who were followed for an average of 7 years, secondary cancers occurred in 26 [7%], a rate which was minimally higher than might be expected and was not associated with any specific therapy, interferon, cladribine or pentostatin).

Sanchez M, Orero M, Marco J, Simó, Linares M, Carbonell F. Fulminant hepatic failure after 2'-deoxycofomycin (pentostatin). *Br J Haematol* 1999; 105: 316. PubMed PMID: 10366248.

(72 year old woman with peripheral T cell lymphoma developed jaundice 6 weeks after starting weekly infusions of pentostatin [bilirubin 24.0 mg/dL, ALT 312 U/L, Alk P 2460 U/L], progressing to hepatic failure and death within 2 weeks of onset).

Ho AD, Suciú S, Stryckmans P, De Cataldo F, Willemze R, Thaler J, Peetermans M, et al. Pentostatin in T-cell malignancies--a phase II trial of the EORTC. Leukemia Cooperative Group. *Ann Oncol* 1999; 10: 1493-8. PubMed PMID: 10643542.

(Among 76 patients with T cell malignancies treated with pentostatin, toxicities were mild and self-limited and included liver test abnormalities in 24%, but values above 5 times ULN occurred in only 1 patient).

Ho AD, Suciú S, Stryckmans P, De Cataldo F, Willemze R, Thaler J, Peetermans M, et al. Pentostatin (Nipent) in T-cell malignancies. Leukemia Cooperative Group and the European Organization for Research and Treatment of Cancer. *Semin Oncol* 2000; 27 (2 Suppl 5): 52-7. PubMed PMID: 10877053.

(Same study as described in Ho [1999]).

Tsimberidou AM, Giles F, Duvic M, Fayad L, Kurzrock R. Phase II study of pentostatin in advanced T-cell lymphoid malignancies: update of an M.D. Anderson Cancer Center series. *Cancer* 2004; 100: 342-9. PubMed PMID: 14716770.

(Among 42 adults with T cell malignancies treated with 1 to 12 courses of pentostatin [4 mg/m²], 20 [48%] developed infections, 19% serum creatinine elevations and one Guillain Barre like syndrome; no mention of ALT elevations or hepatotoxicity).

Else M, Ruchlemer R, Osuji N, Del Giudice I, Matutes E, Woodman A, Wotherspoon A, et al. Long remissions in hairy cell leukemia with purine analogs: a report of 219 patients with a median follow-up of 12.5 years. *Cancer* 2005; 104: 2442-8. PubMed PMID: 16245328.

(Among 219 patients with hairy cell leukemia treated with either cladribine [n=34] or pentostatin [n=185], rates of complete remission [81% vs 82%] and 10 year survival [100% vs 96%] were similar).

Cannon T, Mobarek D, Wegge J, Tabbara IA. Hairy cell leukemia: current concepts. *Cancer Invest* 2008; 26: 860-5. PubMed PMID: 18798068.

(Review of the clinical features, course and therapy of hairy cell leukemia; cladribine and pentostatin are first line therapies for this disease and have similar rates of long term response; infections may be less common with pentostatin, but cladribine is given in a single course, whereas pentostatin must be given as multiple courses).

Jain P, Pemmaraju N, Ravandi F. Update on the Biology and Treatment Options for Hairy Cell Leukemia. *Curr Treat Options Oncol* 2014; 15:187-209. PubMed PMID: 24652320.

(Review of the clinical features, etiology and therapy of hairy cell leukemia which has been linked to mutations in the BRAF gene [V600E] and, while usually responsive to therapy with cladribine or pentostatin, promising future approaches include inhibitors of the BRAF signaling pathway).