



Trifluoperazine

Updated: December 16, 2013.

OVERVIEW

Introduction

Trifluoperazine is a phenothiazine and antipsychotic agent that no longer commonly used in clinical practice. Trifluoperazine is a rare cause of clinically apparent acute cholestatic liver injury.

Background

Trifluoperazine (trye" floo oh per' a zeen) is a piperazine phenothiazine derivative which acts by postsynaptic inhibition of dopamine receptors. Trifluoperazine has other peripheral and central nervous system effects, producing both alpha adrenergic stimulation and blocking histamine- and serotonin-mediated effects. Trifluoperazine is indicated for the therapy of acute and chronic psychosis and rarely for nonpsychotic anxiety. Trifluoperazine was approved for use in the United States in 1959 and was formerly a commonly prescribed antipsychotic medication, but has been replaced in recent years in large part by the atypical antipsychotics, which have fewer extrapyramidal side effects. Trifluoperazine is available as tablets of 1, 2, 5 and 10 mg in generic forms and formerly under the brand name of Stelazine. Oral solutions and parenteral formulations are also available. The typical recommended dose in adults in 1 to 2 mg twice daily, increasing as needed to a maximum of 6 mg daily. Common side effects include drowsiness, dizziness, headache, blurred vision, dry mouth, constipation, tremor, restlessness, muscle spasms and weight gain.

Hepatotoxicity

Liver test abnormalities have been reported to occur in a high proportion of patients on long term phenothiazine therapy, but elevations are uncommonly above 3 times the upper limit of normal. The aminotransferase abnormalities are usually mild, asymptomatic and transient, reversing even with continuation of medication. Rare instances of clinically apparent acute liver injury, resembling that due to chlorpromazine, have been reported with trifluoperazine therapy. The onset of jaundice is usually within 1 to 4 weeks, and the pattern of serum enzyme elevations is typically cholestatic or mixed. Immunoallergic features (rash, fever and eosinophilia) were present in some cases but were mild and self-limited; autoantibodies were rare.

Mechanism of Injury

The mechanism by which phenothiazines cause serum aminotransferase elevations is not known. Several features of the clinical presentation of phenothiazine hepatotoxicity (short latency period, fever, eosinophilia) suggest a hypersensitivity reaction, and rechallenge typically causes a rapid recurrence of injury. Trifluoperazine is extensively metabolized by the liver via sulfoxidation and oxidation, and some instances of serum

aminotransferase elevations as well as more clinical apparent liver injury may be caused by production of a toxic intermediate of its metabolism.

Outcome and Management

The serum aminotransferase elevations that occur on trifluoperazine therapy are usually transient and do not require dose modification or discontinuation of therapy. The acute cholestatic hepatitis caused by trifluoperazine is typically self-limited and benign. However, a proportion of cases of phenothiazine cholestatic hepatitis may be followed by prolonged jaundice and cholestasis and vanishing bile duct syndrome. Many patients with chronic cholestasis eventually improve, but they often have persistent enzyme elevations and biliary cirrhosis. No fatalities from trifluoperazine jaundice have been reported. Rechallenge with phenothiazines usually causes a prompt recurrence of the liver injury and should be avoided. Patients with trifluoperazine induced liver injury may have cross sensitivity to other phenothiazines, but generally tolerate haloperidol and the atypical antipsychotics.

Drug Class: [Antipsychotic Agents](#)

Other Drugs in the Subclass, Phenothiazines: [Chlorpromazine](#), [Fluphenazine](#), [Perphenazine](#), [Prochlorperazine](#), [Thioridazine](#)

PRODUCT INFORMATION

REPRESENTATIVE TRADE NAMES

Trifluoperazine – Generic, Stelazine®


DRUG CLASS

Antipsychotic Agents

COMPLETE LABELING

Product labeling at [DailyMed](#), National Library of Medicine, NIH

CHEMICAL FORMULA AND STRUCTURE

DRUG	CAS REGISTRY NUMBER	MOLECULAR FORMULA	STRUCTURE
Trifluoperazine	117-89-5	C ₂₁ -H ₂₄ -F ₃ -N ₃ -S	

ANNOTATED BIBLIOGRAPHY

References updated: 16 December 2013

Zimmerman HJ. Neuroleptic drugs. In, Zimmerman HJ. Hepatotoxicity: the adverse effects of drugs and other chemicals on the liver. 2nd ed. Philadelphia: Lippincott, 1999, pp. 483-91.

(Expert review of hepatotoxicity of neuroleptic drugs including the phenothiazines published in 1999; several hundred cases of chlorpromazine jaundice have been reported due to chlorpromazine, usually cholestatic, arising after 1-5 weeks, often with fever and eosinophilia, sometimes causing vanishing bile duct syndrome; other phenothiazines have only rarely been linked to liver injury, except for prochlorperazine).

Larrey D, Ripault M-P. Phenothiazines. Hepatotoxicity of psychotropic drugs and drugs of abuse. In, Kaplowitz N, DeLeve LD, eds. Drug-induced liver disease. 3rd ed. Amsterdam: Elsevier, 2013, p. 452.

(Review of phenothiazine hepatotoxicity mentions that liver enzyme elevations arise in up to 40% of patients taking phenothiazines and hundreds of cases of chlorpromazine jaundice have been published, frequency ~0.5-1%; onset in 2-5 weeks, usually acute cholestatic hepatitis with jaundice and pruritus; prodrome of fever and abdominal pain is common; prolonged course in 7% but usually benign; other phenothiazines have been linked to liver injury similar to that of chlorpromazine, "but with a lower frequency").

Meyer JM. Pharmacotherapy of psychosis and mania. In, Brunton LL, Chabner BA, Knollman BA, eds. Goodman & Gilman's the pharmacological basis of therapeutics. 12th ed. New York: McGraw-Hill, 2011, pp. 417-56.

(Textbook of pharmacology and therapeutics).

Kohn N, Myerson RM. Cholestatic hepatitis associated with trifluoperazine. N Engl J Med 1961; 264: 549-50. PubMed PMID: 13757600.

(48 year old man developed jaundice and abdominal pain 3 weeks after starting trifluoperazine [bilirubin 5.2 mg/dL, AST 95 U/L, Alk P ~10 x ULN, eosinophils 6%], resolving within 4 weeks of stopping).

McQueen EG. Toxic effects of phenothiazine tranquilizers. N Z Med J 1963; 62: 460-2. PubMed PMID: 14073060.

(Review of the phenothiazines and their side effects: "Jaundice has occurred in about 1% of patients taking chlorpromazine, and also, although less frequently, in patients taking one of the more recently developed analogues").

Walker CO, Combes B. Biliary cirrhosis induced by chlorpromazine. Gastroenterology 1966; 51: 631-40. PubMed PMID: 5926937.

(32 year old woman and 31 year old man developed persistent jaundice [>4 years], cholestasis and liver fibrosis 3 and 4 weeks after starting chlorpromazine; acute cholestatic hepatitis evolving into chronic form with biopsies showing cirrhosis and complications of portal hypertension, most likely representing vanishing bile duct syndrome).

Margulies AI, Berris B. Jaundice associated with the administration of trifluoperazine. Can Med Assoc J 1968; 98: 1063-4. PubMed PMID: 5652451.

(26 year old woman developed fever, rash and pruritus 2 weeks after starting trifluoperazine [bilirubin 6.4 mg/dL, ALT 169 U/L, Alk P ~1.8 times normal, 8% eosinophils], resolving within 1 month stopping and later tolerating haloperidol without recurrence).

Ishak KG, Irey NS. Hepatic injury associated with the phenothiazines. Clinicopathologic and follow-up study of 36 patients. Arch Pathol 1972; 93: 283-304. PubMed PMID: 5017281.

(Review of 36 biopsies from patients with phenothiazine induced liver injury from the files of the Armed Forces Institute of Pathology, 33 chlorpromazine, 3 prochlorperazine; mean onset 15 days after starting; eosinophilia in

73%, mean bilirubin 12.4 mg/dL, Alk P ~8 fold elevated, ALT 146 U/L; 6 [17%] had prolonged course for 10-16 months).

Døssing M, Andreassen PB. Drug-induced liver disease in Denmark. An analysis of 572 cases of hepatotoxicity reported to the Danish Board of Adverse Reactions to Drugs. *Scand J Gastroenterol* 1982; 17: 205-11. PubMed PMID: 6982502.

(Among 572 cases of drug induced liver disease seen between 1968-78 in Denmark, 51 were attributed to chlorpromazine [9%, ranking 2nd behind halothane], latency averaging 14 days [range 11-46]; 5 deaths; no other phenothiazines mentioned).

Kaplowitz N, Aw TY, Simon FR, Stolz A. Drug-induced hepatotoxicity. *Ann Intern Med* 1986; 104: 826-39. PubMed PMID: 3518564.

(Review of drug induced hepatotoxicity including phenothiazine jaundice).

Munyon WH, Salo R, Briones DF. Cytotoxic effects of neuroleptic drugs. *Psychopharmacology (Berl)* 1987; 91: 182-8. PubMed PMID: 2883697.

(In vitro cell cytotoxicity of 8 neuroleptic drugs were compared; chlorpromazine was more toxic than haloperidol or loxapine, but similar to other phenothiazines).

Regal RE, Bili JE, Glazer HM. Phenothiazine-induced cholestatic jaundice. *Clinical Pharmacy* 1987; 6: 787-94. PubMed PMID: 2905941.

(Review of phenothiazine induced liver injury; cross sensitivity is rare "but does exist").

Pillans PI. Drug associated hepatic reactions in New Zealand: 21 years experience. *NZ Med J* 1996; 109: 315-9. PubMed PMID: 8816722.

(Over 21 year period in New Zealand, there were 943 reports of liver injury involving 205 drugs; chlorpromazine was in the top 20 drugs implicated accounting for 2.7% of cases; prochlorperazine was cause of 4 cases, but other phenothiazines not mentioned).

Allison DB, Mentore JL, Heo M, Chandler LP, Cappelleri JC, Infante MC, Weiden PJ. Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am J Psychiatry* 1999; 156: 1686-96. PubMed PMID: 10553730.

(Systematic review of 81 articles on weight change with antipsychotics; using change after 10 weeks to compare agents: clozapine +5.7, olanzapine +4.2, chlorpromazine +4.2, risperidone +1.7, loxapine +0.6, haloperidol +0.5, ziprasidone +0.3, molindone -0.1, and pimozide -2.7 kilograms).

Sabaté M, Ibáñez L, Pérez E, Vidal X, Buti M, Xiol X, Mas A, et al. Risk of acute liver injury associated with the use of drugs: a multicentre population survey. *Aliment Pharmacol Ther* 2007; 25: 1401-9. PubMed PMID: 17539979.

(Among 126 cases of drug induced liver injury seen in Spain between 1993 and 2000, 3 were due to chlorpromazine with relative risk of 613: frequency of 261 per 100,000 person year exposures; other phenothiazines not mentioned).

Chalasanani N, Fontana RJ, Bonkovsky HL, Watkins PB, Davern T, Serrano J, Yang H, Rochon J; Drug Induced Liver Injury Network (DILIN). Causes, clinical features, and outcomes from a prospective study of drug-induced liver injury in the United States. *Gastroenterology* 2008; 135: 1924-34. PubMed PMID: 18955056.

(Among 300 cases of drug induced liver disease in the US collected between 2004 and 2008, none were due to phenothiazines).

Flanagan RJ. Fatal toxicity of drugs used in psychiatry. *Hum Psychopharmacol* 2008; 23 Suppl 1: 43-51. PubMed PMID: 18098225.

(Deaths from fatal poisonings decreased in England and Wales between 1993-2004, antipsychotic overdose fatalities higher for phenothiazines than atypicals; deaths/million prescriptions being 29 for chlorpromazine, 15.5 thioridazine, 3.9 trifluoperazine, 13.3 olanzapine, 21 clozapine and 31.3 quetiapine).

Reuben A, Koch DG, Lee WM; Acute Liver Failure Study Group. Drug-induced acute liver failure: results of a U.S. multicenter, prospective study. *Hepatology* 2010; 52: 2065-76. PubMed PMID: 20949552.

(Among 1198 patients with acute liver failure enrolled in a US prospective study between 1998 and 2007, 133 were attributed to drug induced liver injury including 4 due to psychotropic agents; one each for quetiapine, nefazodone, fluoxetine and venlafaxine, but none for phenothiazines).

Sharifi V, Amini S, Ghaeli P, Saeedinia A. Trifluoperazine-induced cholestatic jaundice. *Iran J Psychiatry* 2010; 5: 117-8. PubMed PMID: 22952504.

(44 year old man developed jaundice 4 weeks after starting trifluoperazine [bilirubin 3.9 mg/dL, ALT 235 U/L, Alk P 744 U/L], resolving with stopping and not recurring after switching to risperidone).

Molleston JP, Fontana RJ, Lopez MJ, Kleiner DE, Gu J, Chalasani N; Drug-induced Liver Injury Network. Characteristics of idiosyncratic drug-induced liver injury in children: results from the DILIN prospective study. *J Pediatr Gastroenterol Nutr* 2011; 53: 182-9. PubMed PMID: 21788760.

(Among 30 children with suspected drug induced liver injury, half [n=15] were due to antimicrobials [minocycline 4, INH 3, azithromycin 3] and the rest largely due to CNS agents and anticonvulsants; one case was attributed to perphenazine).